

University of Groningen

Identification of distinct nisin leader peptide regions that determine interactions with the modification enzymes NisB and NisC

Khusainov, Rustem; Moll, Gert N.; Kuipers, Oscar P.

Published in:
FEBS Open Bio

DOI:
[10.1016/j.fob.2013.05.001](https://doi.org/10.1016/j.fob.2013.05.001)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2013

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Khusainov, R., Moll, G. N., & Kuipers, O. P. (2013). Identification of distinct nisin leader peptide regions that determine interactions with the modification enzymes NisB and NisC. *FEBS Open Bio*, 3, 237-242.
<https://doi.org/10.1016/j.fob.2013.05.001>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supplementary material

Identification of distinct nisin leader peptide regions that determine interactions with the modification enzymes NisB and NisC


Rustem Khusainov¹, Gert N. Moll² and Oscar P. Kuipers^{1,3*}

¹ Molecular Genetics Dept., University of Groningen, Nijenborgh 7, 9747 AG Groningen, the Netherlands

² Lanthio Pharma, PO Box 719, 9700 AS Groningen, the Netherlands

³ Kluyver Centre for Genomics of Industrial Fermentation, 9747 AG Groningen, the Netherlands

Figure S1. The HMM logos show the probability of each amino acid (height of the letter) and are scaled (width of the letter) according to how many sequences contributed to that position (i.e. narrow letters were generated from a smaller number of sequences than wider letters). The HMM logos were generated from 12 class I lantibiotic leader peptide sequences.



Accession numbers of the class I lantipeptides used for the generation of HMM-logos:

NisA P13068

NisZ P29559

Nisin Q BAD05046

Nisin F ABU45463

Nisin U Q2QBT0

Epilancin K7 AAA79236

Subtilin P10946

Pep5 CAA90023

Ericin S AAL15569

Ericin A AAL15567

Epidermin P08136

Gallidermin P21838

Figure S2. Matrix-assisted laser desorption ionization–time-of-flight (MALDI-TOF) mass spectra of nisin variants modified *in vivo* by NisBTC enzymes in *L.lactis* and purified by Ni-NTA. –Met stands for nisin without the initial methionine, +Met stands for nisin with the initial methionine. x denotes the number of dehydrations.

